# S•Tag<sup>™</sup>: A Multipurpose Fusion Peptide for Recombinant Proteins

Mark McCormick and Robert Mierendorf - Novagen, Inc.

he S•Tag sequence is a novel fusion peptide tag for recombinant proteins that allows detection by a rapid, sensitive homogeneous assay or by colorimetric detection in Western blots. Proteins can also be rapidly purified from crude extracts using a one step affinity separation method. The system is based on the strong interaction between the 15aa S•Tag and 103aa S-protein, both of which are derived from RNase A, resulting in reconstituted ribonucleolytic activity (RNase S; see sidebar article). This report demonstrates the utility of the S•Tag sequence for quantitative assay, blot detection and purification of recombinant proteins.

## S•Tag Vectors

The S•Tag peptide sequence, LysGluThrAlaAlaAlaLysPheGluArgGlnHisMetAspSer is encoded by pCITE<sup>TM</sup>-3 and -4 series for expression of target proteins by *in vitro* translation, and by pET-29 and -30 vector series for expression in *E. coli*. In all of these vectors the S•Tag fusion is at the N-terminus of the target protein, and each series carries a multiple cloning site region available in all three reading frames. Other useful features such as protease cleavage sites and His•Tag<sup>®</sup> sequences are also included (see maps on pp. 8–9).

The pCITE vectors encode the EMCVderived Cap-Independent Translation Enhancer sequence (CITE) and are designed for optimal production of proteins *in vitro* (1). The CITE sequence acts as an internal entry point for initiation of translation by eukaryotic ribosomes and enhances the translation efficiency of proteins as much as 16-fold in reticulocyte lysates, such as Novagen's Red Nova<sup>®</sup> Lysate and Single Tube Protein<sup>TM</sup> System.

The pET-29 and pET-30 vector series are the newest members of the pET System, featuring high-level T7 promoter-driven protein expression in *E. coli*. Experiments at Novagen have indicated that the S•Tag sequence is efficiently expressed in *E. coli* and has not negatively affected the expression level of a variety of proteins when fused to the N-terminus (data not shown). The presence of an S•Tag or His•Tag sequence allows proteins expressed from pET-29 and pET-30 to be rapidly purified using either the S•Tag Purification Kit or the His•Bind<sup>®</sup> resin. Extremely pure, intact target proteins can be prepared using a combination of Nterminal S•Tag and C-terminal His•Tag sequences in a sequential dual affinity purification strategy (2).

Site-specific protease cleavage sites encoded by the vectors allow removal of the leader sequence without significant secondary cleavage. Novagen's unique Biotinylated Thrombin is convenient for this purpose because the enzyme can easily be removed with streptavidin-agarose after the reaction is complete. The pCITE and pET S•Tag vectors are also available as



## T-vectors prepared for direct, reliable cloning of PCR amplified DNA.



Figure 2. Induction time course analysis of two target proteins cloned in pET-30b(+). Whole cell extracts were prepared by removing aliquots from the culture at the indicated times after the addition of IPTG to induce expres-

Whole cell extracts were prepared by removing aliquots from the culture at the indicated times after the addition of IPTG to induce expression. The left-hand gels were stained with Coomassie blue; the right-hand gels were loaded with a 1/100 dilution of the same samples, transferred to ntirocellulose and stained with S•protein-alkaline phosphatase conjugate followed by color development with NBT/BCIP. The target proteins and host strains were  $\beta$ -galactosidase and NovaBlue(DE3) (top), and a recombinant antibody and BL21(DE3) (bottom).

## S•Tag Rapid Assay

The S•Tag rapid assay is based on the reconstitution of ribonucleolytic (RNase S) activity. A sample containing the target protein is added to a buffer containing purified S-protein and the ribonuclease substrate poly(C). After a brief incubation the reaction is stopped with trichloroacetic acid and the resulting precipitate removed by centrifugation. Activity is measured by reading the absorbance of the supernatant at 280nm, which increases as the poly(C) is broken down into acid-soluble nucleotides by the enzyme (3). In the presence of excess S-protein the level of RNase S activity is directly proportional to the amount of recombinant protein present in the sample.

Typical assay profiles are shown in Figure 1. A linear signal was obtained during a 5 minute incubation at 37°C. In this experiment,  $2\mu$ l of crude translation mix and  $2\mu$ l of a 1/100 dilution of *E. coli* lysed with 1% SDS were used as samples. Appropriate blanks serve as controls for any endogenous RNase activity, which appears to be minimal under the conditions of the Rapid Assay. By comparing the absorbance profile of a known standard (provided in the kit), the molar concentration of the S•Tag target protein can be calculated.

Unlike amino acid incorporation assays (both radioactive and non-radioactive), S•Tag Rapid Assay is independent of protein size, amino acid composition and endogenous amino acid pool size. It, therefore, represents a much more accurate method for measuring in vitro translation efficiencies. The method also is extremely versatile for measuring protein expression in cells, since it can be applied to both soluble and insoluble proteins. Up to  $10\mu$ l of a 1/100dilution of 6M urea or guanidine HCl can be added to the assay with little effect (data not shown). Multiple samples can easily be screened for expression levels by preparing crude extracts of whole cells in 1% SDS. Since the assay will detect as little as 20fmol target protein in a 5 minute incubation, even poorly expressed proteins can be measured with a high degree of accuracy. Typically, 1ml of induced culture provides enough material for over 1000 S•Tag Rapid Assays or 100 S•Tag Western Blots.

#### S•Tag Western Blot

Western blotting of S•Tag proteins provides a second means of detection and allows visualization of protein integrity. The S•Tag Western Blot Kit is based on the interaction between the S•Tag sequence and S-protein:biotin or enzyme conjugates. Colorimetric substrates are included which allow as little as 250pg of target protein to be visualized using a 30 minute protocol. Figure 2 shows a time course of induction of  $\beta$ -galactosidase and a recombinant antibody cloned in pET-30b(+). The S•Tag western blot shows high specificity for target proteins (and their amino-terminal-containing breakdown products) with very low background staining of other *E. coli* proteins. The gels also contain Novagen's Perfect

Protein<sup>TM</sup> markers, a set of 7 proteins with precise molecular weights at convenient intervals (15, 25, 35, 50, 75, 100, and 150kd). The markers are known amounts of defined recombinant proteins containing the S•Tag and thus serve as precise internal standards for S•Tag western blots.

## S•Tag Affinity Purification

The high affinity interaction between S-protein and S•Tag also can be applied to purification of target proteins. The S•Tag Purification Kit contains S-protein immobilized on agarose beads to achieve rapid, *continued on page 6* 

## S-Peptide in Protein Fusion Systems: History and Significance

Jin-Soo Kim and Ronald T. Raines — Department of Biochemistry, University of Wisconsin–Madison

RNA: Almost forty years ago, Fred Richards and coworkers discovered that the protease subtilisin prefers to cleave a single peptide bond in native RNA: Almost (1). The product of this cleavage, ribonuclease S (RNA: S), consists of two tightly-associated fragments: S-peptide (residues 1–20) and S-protein (residues 21–124). Although neither fragment alone has any ribonuclease activity, RNA: S has enzymatic activity similar to that of intact RNA: A Richards and Harold Wyckoff determined the structure of crystalline RNA: S has a single peptide of the structure of crystalline crystalline RNA: S has a single peptide of the structure of crystalline crystalline crys

The S-peptide fragment of RNase A has played an important role in the history of biochemistry. Before molecular biologists were able to use recombinant DNA technology to explore protein structure-function relationships, organic chemists synthesized analogs of S-peptide and studied their complexes with S-protein. These studies provided much information on the role of individual residues in RNase S. Most significantly, Chris Anfinsen and coworkers found that only residues 1–15 of S-peptide were necessary to form a fully functional complex with S-protein (3). This shorter fragment is called "S15" (or "S•Tag<sup>™</sup>"</sup> by Novagen) and its complex with S-protein is called "RNase S."

The detection, immobilization, and purification of proteins is idiosyncratic and can be problematic. Fortunately, these processes can be generalized by using recombinant DNA technology to produce fusion proteins in which target proteins are fused to carrier polypeptides (4). The affinity of the carrier for a specific ligand enables the facile detection, immobilization *(continued on page 6)* 

## ARTICLE

## continued from page 5

one-step purification of up to 1mg of target protein from crude extracts. Several purification strategies are available depending on the application. For purification of soluble proteins under native conditions, target proteins bound to the matrix can be eluted by cleavage with biotinylated thrombin, leaving the S•Tag peptide behind. The biotinylated thrombin is then quantitatively removed with streptavidin-agarose. Figure 3 shows an example of the purification achieved under these conditions.

If retention of the S•Tag fusion is desired, proteins can be eluted with 2M sodium thiocyanate. Target proteins found in inclusion bodies can also be purified by solubilization in 6M urea followed by column binding and washing in 2M urea and elution with thiocyanate

The S•Tag purification strategy is recommended for small to medium scale applications. For producing target proteins on a larger scale, a more economical strategy takes advantage of the His•Tag<sup>®</sup> fusion peptide also encoded by these vectors for His•Bind<sup>®</sup> metal chelation chromatography. His•Tag proteins can be purified under a wide range of either native or denaturing conditions. The His•Bind affinity resin has a very high capacity for target proteins; up to 20mg of fusion protein can be bound on a single 2.5ml column. The method is based

## (continued from page 5)

and purification of a fusion protein. The most sensitive detection methods rely on the catalytic activity of an enzymic carrier. Unfortunately, enzymes are relatively large, and are, therefore, more likely than simple peptides to perturb a target protein or to be immunogenic.

The wealth of information that has been accumulated on RNase S suggested to us that this non-covalent complex may provide a useful carrier and ligand for a fusion protein system. We have drawn on this wealth to develop a fusion protein system in which S•Tag is the carrier and S-protein is the ligand (5). The S-Tag carrier combines a small size (15 amino acid residues) with a high sensitivity of detection (10ng of a typical fusion protein in solution or 1pg in a polyacrylamide gel, both without antibodies or radioactivity). S•Tag has several additional properties that are desirable in a carrier. For example, S-Tag [which is composed of 7 charged polar, 3 uncharged polar, and 5 nonpolar residues] is an excessively soluble peptide with little structure and net charge at neutral pH. The S•Tag carrier is therefore unlikely to interfere with the proper folding or function of a fused target protein. Also, the topology of RNase S is such that target proteins fused to either terminus of S-Tag allow binding to S-protein. Finally, the affinity between S-Tag and S-protein can be finetuned by rational mutagenesis (5). Together, these properties make S•Tag an extremely useful and versatile carrier in protein fusion systems.

3. Potts, J.T., Jr., Young, D.M. and Anfinsen, C.B. (1963) J. Biol. Chem. 238, 2593-2594.

1 2 3 4 - β-gal - HSA - thrombin

#### Figure 3. S•Tag purification.

S • Tag  $\beta$ -galactosidase from pET-29b(+) was purified from a crude soluble fraction under native conditions. Samples were analyzed by SDS-PAGE and Coomassie blue staining.

- 1. Crude extract
- 2. Protein released with Biotinylated Thrombin
- 3. Thrombin standard (HSA added as stabilizer)
- 4. Protein after biotinylated thrombin removal

on the interaction between immobilized nickel cations and the His•Tag oligohistidine domain, and elution is achieved with imidazole. In practice, this strategy produces an equivalent amount of pure protein for usually less than 1/20 the cost of affinity methods based on immobilized protein ligands such as antibodies or avidin.

The availability of two independent affinity strategies (S•Tag and His•Tag) also allows a dual affinity approach to purify target proteins "from both ends," effectively eliminating all non-full length products and producing exquisitely pure preparations (2).

## Summary

The S•Tag System provides a unique set of research tools suitable to quantification, detection and purification of expressed proteins. Perhaps its most unique feature is the reconstitution of enzymic activity that is easily assayed with high sensitivity. The small size, low cost and specificity of the S-protein may provide additional advantages over other enzyme-based detection methods in a variety of applications, which we continue to investigate.

### References

- Parks, G.D., Duke, G.M. and Palmenberg, A.C. (1986) *J. Virol.* 60, 376-384; Duke, G.M., Hoffman, M.A. and Palmenberg, A.C. (1992) *J. Virol.* 66, 1602-1609.
- 2. Kim, J.-S. and Raines, R.T. (1994) *Anal. Biochem.***219**, 165-166.
- 3. Zimmerman, S.B. and Sundeen, G. (1965) *Anal. Biochem.* **10**, 444-449.

Richards, F.M. (1955) Compt. Rend. Trav. Lab Carlsberg, Ser. Chim. 29, 322-328; Richards, F.M. and Vithayathil, P.J. (1959) J. Biol. Chem. 234, 1459-1465; Richards, F.M. (1992) Protein Sci. 1, 1721-1730.

Wyckoff, H.W., Hardman, K.D., Allewell, N.M., Inagami, T., Tsernoglou, D., Johnson, L.N. and Richards, F.M. (1967a) *J. Biol. Chem.* 242, 3749-3753; Wyckoff, H.W., Hardman, K.D., Allewell, N.M., Inagami, T., Johnson, L.N. and Richards, F.M. (1967b). *J. Biol. Chem.* 242, 3984-3988.

For recent reviews, see: Uhlén, M. and Moks, T. (1990) *Meth. Enzymol.* **185**, 129-143; Ford, C.F., Suominen, I. and Glatz, C.E. (1991) *Prot. Exp. Purif.* **2**, 95-107; Nilsson, B., Forsberg, G., Moks, T., Hartmanis, M. and Uhlén, M. (1992) *Curr. Opin. Struct. Biol.* **2**, 569-575.

<sup>5.</sup> Kim, J.-S. and Raines, R.T. (1993) *Protein Sci.* **2**, 348-356.