

Integrated Solutions for Recombinant Protein Extraction and Stabilization

Reference and Selection Guide

BACTERIAL

MAMMALIAN

PLANT

YEAST

Efficiently Extract Protein

Preserve Protein Function

Prevent Protein Degradation

sigma-aldrich.com/lysis

 **SIGMA**[®]

Protease and Phosphatase Inhibitor Cocktails

sigma-aldrich.com/protinhib

Sample Type	General Use		Bacterial	Mammalian	Plant	Yeast	Histidine-tagged Proteins	Tissue Culture Media	Phosphatase Inhibitors	
									Tyrosine/Alkaline	Serine/Threonine
Catalog Number	P2714	NEW S8820	P8465	P8340	P9599	P8215	P8849	P1860	P5726	P2850
Specificity of Inhibition	Serine, Cysteine, Aspartic & Metalloproteases	Serine, Cysteine, Aspartic, & Metalloproteases	Serine, Cysteine, Aspartic, & Metalloproteases Aminopeptidases	Serine, Cysteine, Aspartic Proteases & Amidopeptidases	Serine, Cysteine, Aspartic and Metalloproteases, Aminopeptidases	Serine, Cysteine, Aspartic and Metalloproteases	Serine, Cysteine, Aspartic and Thermolysin-like Proteases & Aminopeptidases	Serine, Cysteine, Aspartic proteases & Aminopeptidases	Acid and Alkaline Phosphatases, as well as Tyrosine Protein Phosphatases	Serine, Threonine Phosphatases and various isozymes of Alkaline Phosphatase
Cocktail Format	Water soluble lyophilized powder	Water or buffer soluble tablet	Lyophilized powder with a vial of DMSO	Solution in DMSO	Solution in DMSO	Solution in DMSO	Solution in DMSO	Solution in DMSO	Solution in water	Solution in DMSO
Usage (amount of sample that 1 ml of cocktail will inhibit) Typically enough for 100 ml of cell lysate, depending on cell type	1 mg of USP pancreatin; typically enough for 100 ml of cell lysate	1 mg of USP pancreatin (1x solution), One tablet generates 100 ml of protease inhibitor solution	20 ml of cell lysate from 4 g (wet weight) of <i>E. coli</i> cells	100 ml of cell lysate from 20 g (wet weight) of bovine liver or 10 ml of cell lysate from CHO cells at 10 ⁸ cells per ml	100 ml of cell lysate from 30 g (wet weight) of plant tissue	100 ml of cell lysate from 20 g (wet weight) of <i>Saccharomyces cerevisiae</i> cells	100 ml of cell lysate from 20 g (wet weight) of <i>E. coli</i> cells or 10 g (wet weight) of baculovirus infected cells	Dilution of ≥1:200 in tissue culture media to prevent proteolytic degradation of secreted proteins	100,000 x g supernatant from human placenta, bovine liver, rabbit muscle, A431 or Jurkat cell extracts at a protein concentration of approximately 5 mg/ml	
Cocktail Components (and concentrations when available)	AEBSF, 2 mM Aprotinin, 0.3 μM Bestatin, 130 μM EDTA, 1 mM E-64, 14 μM Leupeptin, 1 μM	AEBSF, 2 mM Aprotinin, 0.3 μM Bestatin, 130 μM EDTA, 1 mM E-64, 14 μM Leupeptin, 1 μM	AEBSF, 23 mM Bestatin, 2 mM EDTA, 100 mM E-64, 0.3 mM Pepstatin A, 0.3 mM	AEBSF, 104 mM Aprotinin, 80 μM Bestatin, 4 mM E-64, 1.4 mM Leupeptin, 2 mM Pepstatin A, 1.5 mM	AEBSF Bestatin E-64 Leupeptin Pepstatin A 1,10-Phenanthroline	AEBSF, 100 mM E-64, 1.4 mM Pepstatin A, 2.2 mM 1,10-Phenanthroline, 500 mM	AEBSF Bestatin E-64 Pepstatin A Phosphoramidon	Aprotinin Bestatin E-64 Leupeptin Pepstatin A	Imidazole Sodium Molybdate Sodium orthovanadate Sodium Tartrate	Bromotetramisole Cantharidin Microcystin LR
Additional Special / Specific Applications (Tested on but not limited to)	Optimized and tested for general use	Optimized and tested for general use	Various <i>E. coli</i> strains	Use at 1:100 for undiluted blood samples, liver tissue extracts, CHO cell lysates	Tested on extracts from kidney bean, pea, wheat, tobacco, and <i>Arabidopsis</i>	<i>Saccharomyces cerevisiae</i> cells	<i>E. coli</i> cells; baculovirus-infected <i>Spodoptera frugiperda</i> pupal ovary cells	After 48 hours in medium, add fresh inhibitors in fresh medium	Tested on various animal tissues (bovine liver, human placenta, rabbit muscle) and A431 and Jurkat cell lines	Tested on various animal tissues (bovine liver, human placenta, rabbit muscle) and A431 and Jurkat cell lines
Storage and Stability (unopened)	-20 °C ≥ 4 years	2-8 °C ≥ 4 years	-20 °C ≥ 4 years	At least 4 years at -20 °C; 8 months at 2-8 °C; and 2 months at RT	At least 4 years at -20 °C; 8 months at 2-8 °C; and 2 months at RT	At least 4 years at -20 °C; 8 months at 2-8 °C; and 2 months at RT	At least 4 years at -20 °C; 3 months at 2-8 °C; and 2 weeks at RT	At least 4 years at -20 °C; 8 months at 2-8 °C; and 2 months at RT	2-8 °C ≥ 2 years	2-8 °C ≥ 4 years

Coming Soon! SigmaFAST™ Protease Inhibitor Tablet for General Use (Cat. No. S8820)

Protease and Phosphatase Inhibitors *Available Individually*

Protease Inhibitors	Cat. No.	Application
AEBSF	A8456	Inhibits serine proteases, such as trypsin and chymotrypsin
Aprotinin saline solution Aprotinin, affinity purified	A6279 A4529	Inhibits serine proteases such as trypsin, chymotrypsin, plasmin, trypsinogen, urokinase, and kallikrein. Also inhibits human leukocyte elastase, but not pancreatic elastase
Bestatin, hydrochloride	B8385	Inhibits aminopeptidases, such as leucine aminopeptidase proteases
Cantharidin	C7632	Inhibits protein phosphatase 2A (PP-2A)
Chymostatin	C7268	Inhibits serine and cysteine proteases
E-64	E3132	Inhibits cysteine proteases such as calpain, papain, and cathepsin B
EDTA disodium salt dihydrate EDTA, 0.5 M solution disodium salt	E5134 E7889	Inhibits metalloproteases; chelates; permeabilizes Gram-negative bacteria
EGTA	E8145	Inhibits metalloproteases; chelates
Imidazole	I0125	Inhibits alkaline phosphatases
Leupeptin hemisulfate Leupeptin, TFA salt Leupeptin hydrochloride	L2884 L2023 L9783	Inhibits both serine and cysteine proteases such as calpain, trypsin, papain, and cathepsin B
Microcystin LR	M2912	Inhibits protein phosphatases 1 and 2A (PP-1 and PP-2A)
Pepstatin A Pepstatin A, 90% pure	P4265 P5318	Inhibits acid proteases such as pepsin (human or porcine), renin, cathepsin D, chymosin (bovine rennin), and protease B
1,10-Phenanthroline	P9375	Inhibits metalloproteases
Phosphoramidon	R7385	Inhibits thermolysin and collagenase
PMSF	P7626	Inhibits serine proteases
Sodium molybdate	M1003	Inhibits acid and phosphoprotein phosphatases
Sodium orthovanadate	S6508	Inhibits a number of ATPases, protein tyrosine phosphatases, and other phosphate-transferring enzymes
Sodium tartrate	S4797	Inhibits alkaline phosphatases

Organism


BACTERIA

WHOLE CELL DETERGENT LYSIS

WHOLE

Product	CellLytic™ B	CellLytic B 2x	CellLytic B 10x	CellLytic Express
Catalog number	B7435	B7310	C8740	C1990
Description	A ready-to-use solution containing 40 mM Tris-HCl, pH 8, and a proprietary, non-denaturing detergent formulation of zwitterionic detergents	A more concentrated (2x), ready-to-use solution containing 40 mM Tris-HCl, pH 8, and a proprietary, non-denaturing formulation detergent	A highly concentrated (10x) solution containing a proprietary, non-denaturing detergent formulation and no buffering components	A ready-to-use, all-in-one, non-denaturing proprietary powder formulation of detergent and enzymes optimized for one-step extraction of proteins from bacterial cells (in-culture cell lysis)
Package sizes	50 ml, 500 ml (Sufficient for processing 5 g and 50 g of wet cell paste, respectively)	50 ml, 250 ml (Sufficient for processing up to 10 g and 50 g of wet cell paste, respectively)	10 ml, 50 ml, 100 ml (Sufficient for processing ~10 g, 50 g, and 100 g of wet cell paste, respectively)	25 ml (1.26 g) 10 x 25 ml 6 x 500 ml (25.2 g)
Sample type	<ul style="list-style-type: none"> Bacterial cell pellets (lysis and protein extraction) Isolation of inclusion bodies 	<ul style="list-style-type: none"> Bacterial cell pellets (concentrated at 2x for more difficult to lyse organisms) Isolation of inclusion bodies 	<ul style="list-style-type: none"> Bacterial cell pellets (concentrated at 10x in water; dilute for use into buffer of choice) Isolation of inclusion bodies 	<ul style="list-style-type: none"> Direct lysis of bacterial cell cultures (without centrifugation) Affinity purification of proteins without the need for special equipment
Strain/Cell line compatibility	<ul style="list-style-type: none"> Optimized for <i>E. coli</i> strain BL21 DH5α JM109 Other similar cells 	<ul style="list-style-type: none"> Optimized for <i>E. coli</i> strain BL21 DH5α JM109 Other similar cells 	<ul style="list-style-type: none"> Optimized for <i>E. coli</i> strain BL21 DH5α JM109 Other similar cells 	<ul style="list-style-type: none"> Optimized for <i>E. coli</i> strain BL21 DH5α JM109 Other similar cells
Usage	<u>Small scale:</u> 0.4 ml of CellLytic B per 1.5 ml culture pellet <u>Large scale:</u> 10-20 ml per g of wet cell paste	<u>Small scale:</u> 0.2 ml of CellLytic B 2x per 1.5 ml of culture pellet <u>Large scale:</u> 5 ml per g of wet cell paste	Test 5, 10, and 20-fold dilutions into the desired buffer; add diluted solution at 10-20 ml per g of wet cell paste	<u>Small scale:</u> one 25 ml bottle per 25 ml of culture <u>Large Scale:</u> one 500 ml bottle for 500 ml of culture *For cultures <25 ml, add 0.05 g of powder per ml of culture
Type of detergent/ Dialyzable?	Zwitterionic/Yes	Zwitterionic/Yes	Zwitterionic/Yes	Zwitterionic/Yes
Protein active?	Yes – Suitable for protein activity assays	Yes – Suitable for protein activity assays	Yes – Suitable for protein activity assays	N/A; typically resin-bound protein-activity dependent on isolation/elution methods
Compatible purification and detection products	<ul style="list-style-type: none"> Antibody/EZview™ Affinity Gels FLAG® HIS-Select™/IMAC GST Protease Inhibitor Cocktails Lysozyme solution (L3790) 	<ul style="list-style-type: none"> Antibody/EZview Affinity Gels FLAG HIS-Select/IMAC GST Protease Inhibitor Cocktails Lysozyme solution (L3790) 	<ul style="list-style-type: none"> Antibody/EZview Affinity Gels FLAG HIS-Select/IMAC GST Protease Inhibitor Cocktails Lysozyme solution (L3790) 	<ul style="list-style-type: none"> Antibody/ EZview™ Affinity Gels FLAG® HIS-Select™/IMAC GST Protease Inhibitor Cocktails Lysozyme solution (L3790)
Quantitation and staining capabilities	<ul style="list-style-type: none"> Protein assay Protein staining 	<ul style="list-style-type: none"> Protein assay Protein staining 	<ul style="list-style-type: none"> Protein assay Protein staining 	<ul style="list-style-type: none"> Protein assay Protein staining
Applications	<ul style="list-style-type: none"> Affinity purification SDS-PAGE Western blotting Dot blotting Immunoprecipitation 	<ul style="list-style-type: none"> Affinity purification SDS-PAGE Western blotting Dot blotting Immunoprecipitation 	<ul style="list-style-type: none"> Affinity purification SDS-PAGE Western blotting Dot blotting Immunoprecipitation 	<ul style="list-style-type: none"> Affinity purification SDS-PAGE Western blotting Dot blotting Immunoprecipitation
Related products containing this reagent	See CB0050 & CB0500 kits	See B7435 & CB0500 kits	See B7435 & CB0500 kits	CellLytic Express Tablets (C5491)
Physical form	Ready-to-use solution	Concentrated solution (2x)	Concentrated solution (10x)	Easy-to-dissolve powder
Storage and stability	Room temperature 2 years	Room temperature 2 years	Room temperature 2 years	–20 °C 2 years

Argentina

SIGMA-ALDRICH DE ARGENTINA, S.A.
Tel: 54 11 4556 1472
Fax: 54 11 4552 1698

Australia

Sigma-Aldrich Pty., limited
Free Tel: 1800 800 097
Free Fax: 1800 800 096
Tel: 612 9841 0555
Fax: 612 9841 0500

Austria

SIGMA-ALDRICH Handels GmbH
Tel: 43 1 605 81 10
Fax: 43 1 605 81 20

Belgium

Sigma-Aldrich NV/SA.
Free Tel: 0800-14747
Free Fax: 0800-14745
Tel: 03 899 13 01
Fax: 03 899 13 11

Brazil

Sigma-Aldrich Brasil Ltda.
Tel: 55 11 3732-3100
Fax: 55 11 3733-5151

Canada

Sigma-Aldrich Canada Ltd.
Free Tel: 800-565-1400
Free Fax: 800-265-3858
Tel: 905-829-9500
Fax: 905-829-9292

China

SIGMA-ALDRICH CHINA INC.
Tel: 86-21-6386 2766
Fax: 86-21-6386 3966

Czech Republic

SIGMA-ALDRICH s.r.o.
Tel: +420 246 003 200
Fax: +420 246 003 291

Denmark

Sigma-Aldrich Denmark A/S
Tel: 43 56 59 10
Fax: 43 56 59 05

Finland

Sigma-Aldrich Finland
Tel: (09) 350 9250
Fax: (09) 350 92555

France

Sigma-Aldrich Chimie S.à.r.l.
Tel appel gratuit: 0800 211 408
Fax appel gratuit: 0800 031 052

Germany

Sigma-Aldrich Chemie GmbH
Free Tel: 0800-51 55 000
Free Fax: 0800-649 00 00

Greece

SIGMA-ALDRICH (o.m.) Ltd
Tel: 30 210 9948010
Fax: 30 210 9943831

Hungary

SIGMA-ALDRICH Kft
Tel: 06-1-235-9054
Fax: 06-1-269-6470
Ingyenes zöld telefon: 06-80-355-355
Ingyenes zöld fax: 06-80-344-344

India

SIGMA-ALDRICH CHEMICALS
PRIVATE LIMITED
Telephone
Bangalore: 91-80-5112-7272
New Delhi: 91-11-5165 4255
Mumbai: 91-22-2570 2364
Hyderabad: 91-40-5584 5488
Fax
Bangalore: 91-80-5112-7473
New Delhi: 91-11-5165 4266
Mumbai: 91-22-2579 7589
Hyderabad: 91-40-5584 5466

Ireland

Sigma-Aldrich Ireland Ltd.
Free Tel: 1800 200 888
Free Fax: 1800 600 222
Tel: 353 1 4041900
Fax: 353 1 4041910

Israel

Sigma-Aldrich Israel Ltd.
Free Tel: 1-800-70-2222
Tel: 08-948-4100
Fax: 08-948-4200

Italy

SIGMA-ALDRICH S.r.l.
Telefono: 02 33417310
Fax: 02 38010737
Numero Verde: 800-827018

Japan

SIGMA-ALDRICH JAPAN K.K.
Tokyo Tel: 03 5796 7300
Tokyo Fax: 03 5796 7315

Korea

SIGMA-ALDRICH KOREA
Tel: 031-329-9000
Fax: 031-329-9090

Malaysia

SIGMA-ALDRICH (M) Sdn. Bhd
Tel: 603-56353321
Fax: 603-56354116

Mexico

SIGMA-ALDRICH QUÍMICA, S.A. de C.V.
Free Tel: 01-800-007-5300
Free Fax: 01-800-712-9920

The Netherlands

Sigma-Aldrich Chemie BV
Tel Gratis: 0800-0229088
Fax Gratis: 0800-0229089
Tel: 078-6205411
Fax: 078-6205421

New Zealand

SIGMA-ALDRICH PTY., LIMITED
Free Tel: 0800 936 666
Free Fax: 0800 937 777
Tel: 61 2 9841 0500
Fax: 61 2 9841 0500

Norway

SIGMA-ALDRICH NORWAY AS
Tel: 23 17 60 60
Fax: 23 17 60 50

Poland

SIGMA-ALDRICH Sp. z o.o.
Tel: 061 829 01 00
Fax: 061 829 01 20

Portugal

Sigma-Aldrich Química, S.A.
Free Tel: 800 202180
Free Fax: 800 202178
Tel: 21 9242555
Fax: 21 9242610

Russia

SIGMA-ALDRICH RUS, LLC
Tel: +7 (095) 975-1917/3321
Fax: +7 (095) 975-4792

Singapore

Sigma-ALDRICH PTE. LTD.
Tel: 65-67791200
Fax: 65-67791822

South Africa

Sigma-Aldrich
South Africa (Pty) Ltd.
Free Tel: 0800 1100 75
Free Fax: 0800 1100 79
Tel: 27 11 979 1188
Fax: 27 11 979 1119

Spain

SIGMA-ALDRICH QUÍMICA S.A.
Free Tel: 900 101376
Free Fax: 900 102028
Tel: 91 661 99 77
Fax: 91 661 96 42

Sweden

SIGMA-ALDRICH SWEDEN AB
Tel: 020-350510
Fax: 020-352522
Outside Sweden Tel: +46 8 7424200
Outside Sweden Fax: +46 8 7424243

Switzerland

SIGMA-ALDRICH CHEMIE LLC
Swiss Free Call: 0800 80 00 80
Tel: +41 81 755 2828
Fax: +41 81 755 2815

United Kingdom

Sigma-Aldrich Company Ltd.
Free Tel: 0800 717181
Free Fax: 0800 378785
Tel: 01747 833000
Fax: 01747 833313
SAFC (UK): 01202 712305

United States

SIGMA-ALDRICH
P.O. Box 14508
St. Louis, Missouri 63178
Toll-free: 800-325-3010
Call Collect: 314-771-5750
Toll-Free Fax: 800-325-5052
Tel: 314-771-5765
Fax: 314-771-5757

Internet:

sigma-aldrich.com

sigma-aldrich.com

Order/Customer Service 1-800-325-3010 • Fax 1-800-325-5052

Technical Service 1-800-325-5832 • sigma-aldrich.com/techservice

Development/Bulk Manufacturing Inquiries **SAFC**™ 1-800-244-1173

World Headquarters • 3050 Spruce St., St. Louis, MO 63103 • (314) 771-5765

We are committed to the success of our Customers, Employees and Shareholders through leadership in Life Science, High Technology and Service.

The SIGMA-ALDRICH Family



SIGMA



ALDRICH



Fluka



Riedel-de-Haën



SUPELCO



ISOTEC

©2005 Sigma-Aldrich Co. All rights reserved.

SIGMA, SAFC, SAF, SIGMA-ALDRICH, ISOTEC, ALDRICH, FLUKA, and SUPELCO are trademarks belonging to Sigma-Aldrich Co. and its affiliate Sigma-Aldrich Biotechnology LP. Riedel-de-Haën™ trademark under license from Riedel-de-Haën GmbH. TWEEN® is a registered trademark of Uniqema, a business unit of ICI America, Inc. DH5α™ is a trademark of Invitrogen. IGEPAL® is a registered trademark of Rhone-Poulenc AG Co. TRITON™ is a trademark of Union Carbide Corp. Sigma brand products are sold through Sigma-Aldrich, Inc. Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.



IIB
00159-502642
0105



SIGMA-ALDRICH