



New England Biolabs, Inc.
Tel: 800-632-5227 (orders)
Tel: 800-632-7799 (support)
Fax: 978-921-1350
e-mail: info@neb.com
WWW: <http://www.neb.com>

Enterokinase, light chain (enteropeptidase)

#P8070S 0.063 µg \$80 (USA)
#P8070L 0.32 µg \$320 (USA)

Description

Enterokinase is a specific protease that cleaves at Asp-Asp-Asp-Asp-Lys in trypsinogen to produce trypsin. It cleaves after lysine in its preferred cleavage site. It will sometimes cleave at other basic residues, depending on the conformation of the protein substrate. The molecular weight is 26.3 kDa. The apparent molecular weight on SDS-PAGE is 31 kDa.

Source

This preparation is purified from *E. coli* containing a clone of the light chain of the bovine enterokinase gene, fused to a carrier (1,2)

Suggested Reaction Conditions

The amount of enzyme required to cleave a fusion protein in a 16 hour reaction at room temperature ranges from 0.001% to 0.5% (w/w). Cleavage of the above MBP fusion protein under these conditions requires 0.03%.

Concentration

Selling concentration: 5.0 µg/ml

Storage Conditions

20 mM Tris-HCl (pH 7.2 @ 4°C), 200 mM NaCl, 2 mM CaCl₂ and 50% glycerol. Store at -20°C.

References:

1. LaVallie, E.R. et al. (1993) *J. Biol. Chem.* 268, 23311–23317.
2. LaVallie, E.R. and Racie, L., unpublished observations.

THIS PRODUCT IS SOLD UNDER PATENT LICENSE FROM GENETICS INSTITUTE, INC. FOR RESEARCH ONLY. LICENSES DIRECTLY FROM GENETICS INSTITUTE, INC., 87 CAMBRIDGE PARK DRIVE, CAMBRIDGE, MA. 02140.