

Product Contents

ProTEV Protease:

Supplied With:

Cat. #	Size	ProTEV Protease	ProTEV Buffer, 20X	100mM DTT
V6051	1,000 units	V605A	V602A (1ml)	P117B (250µl)
V6052	10,000 units	V605B	V602B (8ml)	P117C (1.25ml)

Description: ProTEV Protease is an engineered form of TEV protease, a highly specific proteolytic enzyme that cleaves within a seven-amino-acid sequence. It can be used to cleave protein fusions that have been engineered to contain the seven-amino-acid sequence at the desired cleavage site. ProTEV Protease also contains an HQ-tag located at the N-terminus of the protein, which allows it to be immobilized on affinity resins and removed from the cleavage reaction.

Formulation: ProTEV Protease is supplied at a concentration of 10u/µl in 50mM Tris-HCl (pH 7.5), 500mM NaCl, 5mM DTT, 1mM EDTA, 50% glycerol, 0.1% Triton® X-100. ProTEV Buffer, 20X, contains 1M HEPES (pH 7.0), 10mM EDTA.

Unit Definition: Four units of ProTEV Protease cleaves ≥85% of 20µg of a test fusion protein in 30 minutes at 30°C.

Storage Conditions: See the Product Information Label for storage recommendations. Avoid multiple freeze-thaw cycles and exposure to frequent temperature changes. See the expiration date on the product label.

Part# 9PIV605

Printed 5/07



Quality Control Assays

Purity: ≥90% pure by SDS-PAGE.

Specificity: In an overdigestion assay (10u ProTEV Protease, overnight at 30°C), ProTEV Protease does not cleave proteins that do not contain the TEV protease sequence.



Promega

Promega Corporation

2800 Woods Hollow Road	
Madison, WI 53711-5399	USA
Telephone	608-274-4330
Toll Free	800-356-9526
Fax	608-277-2516
Internet	www.promega.com

PRODUCT USE LIMITATIONS, WARRANTY DISCLAIMER

Promega manufactures products for a number of intended uses. Please refer to the product label for the intended use statements for specific products. Promega products contain chemicals which may be harmful if misused. Due care should be exercised with all Promega products to prevent direct human contact.

Each Promega product is shipped with documentation stating specifications and other technical information. Promega products are warranted to meet or exceed the stated specifications. Promega's sole obligation and the customer's sole remedy is limited to replacement of products free of charge in the event products fail to perform as warranted. Promega makes no other warranty of any kind whatsoever, and SPECIFICALLY DISCLAIMS AND EXCLUDES ALL OTHER WARRANTIES OF ANY KIND OR NATURE WHATSOEVER, DIRECTLY OR INDIRECTLY, EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, AS TO THE SUITABILITY, PRODUCTIVITY, DURABILITY, FITNESS FOR A PARTICULAR PURPOSE OR USE, MERCHANTABILITY, CONDITION, OR ANY OTHER MATTER WITH RESPECT TO PROMEGA PRODUCTS. In no event shall Promega be liable for claims for any other damages, whether direct, incidental, foreseeable, consequential, or special (including but not limited to loss of use, revenue or profit), whether based upon warranty, contract, tort (including negligence) or strict liability arising in connection with the sale or the failure of Promega products to perform in accordance with the stated specifications.

© 2007 Promega Corporation. All Rights Reserved.

Products may be covered by pending or issued patents or may have certain limitations. Please visit our Web site for more information.

All prices and specifications are subject to change without prior notice.

Product claims are subject to change. Please contact Promega Technical Services or access the Promega online catalog for the most up-to-date information on Promega products.

Part# 9PIV605
Printed in USA 5/07

I. Description

ProTEV Protease is an improved 50kDa version of the Nla protease from tobacco etch virus (TEV) that has been engineered to be more stable than native TEV protease for prolonged enzymatic activity (1–3). TEV protease is a highly site-specific proteolytic enzyme that recognizes the seven-amino-acid sequence EXXXYQ(G/S), most commonly ENLYFQG, with cleavage occurring between glutamine and glycine or serine (4,5). The protease is used to cleave affinity tags from fusion proteins after protein purification. The protease cleaves sequences with a variety of amino acids at the G/S (or P1') position (6). This allows the choice from many different amino acids on the newly formed N-terminus after cleavage. Optimum activity is obtained at pH 7.0 and 30°C, but ProTEV Protease is active over a wide range of pH values (5.5–8.5) and temperatures (4–30°C), allowing a choice of conditions amenable to the protein of interest. ProTEV Protease is easily removed from the cleavage reaction after cleavage using the HQ-tag located at the N-terminus of the protein. ProTEV Protease can also be used to cleave the affinity tag from a fusion protein immobilized on the affinity resin.

II. General Protocol

ProTEV Protease reactions are carried out in 1X ProTEV Buffer plus 1mM DTT. Assemble the following components in a microcentrifuge tube:

Component	Volume
20X ProTEV Buffer, 20X	5µl
100mM DTT	1µl
fusion protein	20µg
ProTEV Protease (10u)	1µl
Water to	100µl

Incubate samples at 30°C. If preferred, the fusion protein can be cleaved at a lower temperature. Remove 20µl aliquots from the digest at 1, 2, 4 and 6 hours. Add an appropriate SDS-PAGE sample buffer to the aliquots, and store at –20°C until ready to analyze. Analyze 10–20µl by SDS-PAGE, and determine percent cleavage by monitoring the disappearance of the full-length fusion protein and appearance of the cleaved products.

Notes:

1. If the fusion protein is too dilute to include 20µg in a 100µl reaction, either decrease the amount of fusion protein cleaved in the assay or increase the volume of the reaction to accommodate the desired amount of fusion protein.
2. ProTEV Protease has a molecular weight of approximately 50kDa on reducing SDS-PAGE gels.

Cleavage of the fusion protein can be optimized by changing the amount of ProTEV Protease added to the reaction, changing the incubation time, and/or changing the incubation temperature of the reaction.

III. Removal of ProTEV Protease from Cleavage Reactions

ProTEV Protease has an HQ-tag at the N-terminus. After cleaving the fusion protein, ProTEV Protease can be removed from the reaction by incubating with metal-affinity resins like MagneHis™ Ni-Particles (Cat.# V8565) or HisLink™ Protein Purification Resin (Cat.# V8823). Follow the binding instructions for the metal-affinity resin of choice. The cleaved protein of interest will be found in the resin flowthrough or supernatant fraction. If ProTEV Protease was used to remove an HQ- or polyhistidine-tag, this polypeptide will also be removed from the reaction.

IV. Cleavage of Fusion Proteins During Affinity Purification

ProTEV Protease can be used to cleave fusion proteins at the end of affinity purification procedures. Rather than eluting the fusion protein, ProTEV Protease will cleave the fusion protein while it is still bound to the resin, leaving the affinity tag bound to the resin and the protein of interest in the column flowthrough. This process will need to be optimized for each fusion protein with respect to the amount of protease used and time required for cleavage. Optimization can begin using the conditions optimized for the digestion of the fusion protein in solution and varying the assay time and temperature as needed.

In general, after binding the fusion protein to the affinity resin and washing according to the manufacturer's instructions, equilibrate the column with 1X ProTEV Buffer plus 1mM DTT. Leave enough of the buffer in the column to allow free movement of the resin while rocking or rotating the column/tube. Add ProTEV Protease to the column. Using the incubation conditions determined during optimization, incubate while rocking or rotating the resin to keep it resuspended in the buffer. The column flowthrough/supernatant will contain the protein of interest without the affinity tag.

Note: More ProTEV Protease may be required for column cleavage than for solution cleavage of the same fusion protein.

V. References

1. Dougherty, W.G. and Parks, T.D. (1991) Post-translational processing of the tobacco etch virus 49-kDa small nuclear inclusion polyprotein: Identification of an internal cleavage site and delimitation of VPg and proteinase domains. *Virology* **183**, 449–56.
2. Carrington, J.C. *et al.* (1993) Internal cleavage and trans-proteolytic activities of the VPg-proteinase (Nla) of tobacco etch potyvirus in vivo. *J. Virol.* **67**, 6995–7000.
3. Kapust, R.B. *et al.* (2001) Tobacco etch virus protease: Mechanism of autolysis and rational design of stable mutants with wild-type catalytic proficiency. *Protein Eng.* **14**, 993–1000.
4. Dougherty, W.G. *et al.* (1989) Characterization of the catalytic residues of the tobacco etch virus 49-kDa proteinase. *Virology* **172**, 302–10.
5. Carrington, J.C. and Dougherty, W.G. (1988) A viral cleavage site cassette: Identification of amino acid sequences required for tobacco etch virus polyprotein processing. *Proc. Natl. Acad. Sci. USA* **85**, 3391–5.
6. Kapust, R.B. *et al.* (2002) The P1' specificity of tobacco etch virus protease. *Biochem. Biophys. Res. Commun.* **294**, 949–55.

VI. Related Products

Product	Size	Cat.#
MagneHis™ Ni-Particles	10ml	V8565
HisLink™ Protein Purification Resin	5ml	V8823