


SYPRO Ruby Protein Gel Stain

Advanced staining technology for 2-D gels and proteomics



- HIGHLY SENSITIVE** Detection limit rivals that of the best silver stains
- SIMPLE** Ready-to-use stain requires no complex fixations or timed steps
- QUANTITATIVE** Broad linear quantitation range and consistent gel-to-gel staining allow for accurate protein expression comparisons
- COMPATIBLE** Staining is compatible with mass spectrometry or microsequencing
- HIGH-THROUGHPUT** Simple staining procedure streamlines processing of multiple gels
- EASY TO VISUALIZE** Can be visualized with UV transilluminators or laser scanners

Created especially for the analysis of proteins in 2-D polyacrylamide gels, SYPRO Ruby protein gel stain is ideal for proteomics. It provides the same low nanogram sensitivity as the best silver staining techniques (Figure 1), but stains more proteins and has a much broader linear quantitation range — extending over three orders of magnitude (Figure 2). These features make it possible to obtain accurate protein quantitation for both highly expressed and minimally expressed proteins in the gel.¹ Using SYPRO Ruby protein gel stain, it is possible to detect as little as a 5% difference in staining intensities with a 95% confidence level.² The bright orange-red-fluorescent signal (~610 nm) has two excitation maxima (~280 and ~450 nm), making it easy to visualize using either a UV light source, such as a standard transilluminator, or a visible light source, such as a laser or a xenon-arc lamp. Accurate quantitation can be achieved using a CCD camera or a laser scanner.

The staining procedure is very straightforward — after a short fixation, gels are simply incubated in the stain and washed (Figure 3). Unlike silver staining, no timed steps are required — gels can be left in the dye solution for long periods without overstaining. This simplified staining protocol results in very consistent gel-to-gel results; same-spot intensity comparisons between identical 2-D gels show a correlation of 0.9.³ In addition, the streamlined staining procedure makes it possible to process multiple gels simultaneously without investing in robotic staining devices. The amount of hazardous organic waste generated is greatly reduced compared to silver staining, minimizing the hassles associated with waste disposal.




Figure 3. Staining with SYPRO Ruby protein gel stain is simple — just fix, stain and wash.

SYPRO Ruby protein gel stain is a noncovalent stain that does not require aldehyde fixation. Thus, it is compatible with subsequent Edman microsequencing or mass spectrometry (Figure 4), either of which can be performed immediately after staining, with no modification steps.^{1,3} As little as 75 fmol of stained protein can be recovered from the gel and accurately identified using MALDI-TOF mass spectrometry.³

Materials Supplied

SYPRO Ruby protein gel stain is supplied as a ready-to-use solution in 200 mL, 1 L and 5 L sizes. The 1 L size stains ~20 minigels or two large 2-D gels. The 5 L size is provided in a box with a convenient spigot for dispensing the stain.

Ordering Information

S-12000	SYPRO Ruby protein gel stain	1 L
S-12001	SYPRO Ruby protein gel stain	200 mL
S-21900	SYPRO Ruby protein gel stain *bulk packaging*	5 L

References

1. Electrophoresis 21, 2509 (2000);
2. Molecular Probes, unpublished data;
3. Electrophoresis 21, 3673 (2000).


For further information contact

MOLECULAR PROBES, INC.

Eugene, Oregon USA
Customer Service: (541) 465-8338
Customer Service Fax: (541) 344-6504
E-mail: order@probes.com
Technical Assistance: (541) 465-8353
Technical Assistance Fax: (541) 465-4593
E-mail: tech@probes.com
For USA and Canada
Toll-Free Order: (800) 438-2209
Toll-Free Order Fax: (800) 438-0228

MOLECULAR PROBES EUROPE BV

Leiden, The Netherlands
Customer Service: +31-71-5236850
Customer Service Fax: +31-71-5233419
E-mail: eurorder@probes.nl
Technical Assistance: +31-71-5233431
Technical Assistance Fax: +31-71-5241883
E-mail: eurotech@probes.nl



www.probes.com

SYPRO is a registered trademark of Molecular Probes, Inc.
 Molecular Imager is a registered trademark of Bio-Rad Laboratories, Inc.
 © Molecular Probes, Inc., 2001.




Figure 1. SYPRO Ruby protein gel stain compared to a silver stain. Proteins from a fibroblast cell lysate were run on identical 2-D gels and stained with SYPRO Ruby protein gel stain (left) or silver stain (right). The grayscale values of the gel stained with SYPRO Ruby dye have been inverted for easier comparison with the silver-stained gel.




Figure 2. Linear quantitation range of SYPRO Ruby protein gel stain compared to silver stain. Amounts of carbonic anhydrase ranging from 1 ng to 1000 ng were separated by SDS-PAGE and stained with SYPRO Ruby protein gel stain. The inset shows the lower part of the range from 1 ng to 60 ng protein. Staining intensities were quantitated using the Bio-Rad Molecular Imager FX System. For comparison, the gray band shows the linear range for the same protein detected with silver staining.




Figure 4. Mass spectroscopic profile of chicken ovalbumin obtained after SDS-PAGE and staining with SYPRO Ruby protein gel stain. Data courtesy of Elena Chernokalskaya and Mary Lopez, Genomic Solutions, Inc.

These products are offered for research purposes only and are not available for resale or other commercial uses without a specific agreement from Molecular Probes, Inc.