

I. Introduction

This protocol is provided for simple, rapid purification of his-tagged proteins in up to 800 µl of clarified lysate from mammalian or bacterial cell samples using the **Capturem His-Tagged Purification Miniprep Kit** (Cat. No. 635710). The columns are suitable for use under native or denaturing conditions, in the presence of additives such as DTT (up to 10 mM), βME (up to 30 mM), TCEP (up to 5 mM), EDTA (up to 10 mM), or glycerol ([see reagent compatibility table for more information](#)).

II. Materials and Reagents

A. Components

- 20 Capturem Miniprep Nickel Columns
- 2 x 15 ml xTractor Buffer
- 10 ml Wash Buffer
- 10 ml Elution Buffer (500 mM NaCl, 20 mM Na₃ PO₄, 500 mM imidazole, pH 7.6)

B. Additional Materials Required

Each purification will require three additional standard 2-ml collection tubes, with or without a cap. These tubes should be used throughout the protocol to collect flowthrough samples that will be saved for SDS-PAGE analysis and/or colorimetric protein assays, e.g., Bradford assays.

III. Sample Preparation

Before beginning the protein purification protocol in Section IV, it is necessary to prepare a cleared cell lysate from your bacterial or mammalian cell pellet. Lysis protocols using the included xTractor Buffer are provided in the [xTractor Buffer & xTractor Buffer Kit User Manual](#). Individual protocols are also available for preparing cell lysates from [bacterial](#), [mammalian](#), [baculovirus](#), and [yeast](#) cultures.

• Bacterial Cell Samples

We recommend starting with a fresh or frozen cell pellet from 2–5 ml of overnight bacterial culture, which should yield 200–800 µl of cleared lysate.

NOTE: When working with bacterial cells, the volume of lysate (containing the overexpressed his-tagged protein of interest) is determined by the amount of wet cell pellet obtained from a starting culture volume of 2–5 ml. For example, a log-phase *E. coli* culture (O.D. = 0.6–0.8), induced for 2–4 hours, would be expected to provide ~20–80 mg of bacterial pellet from 2–5 ml of culture. We recommend adding ~400 µl of xTractor Buffer to each ~20 mg of wet bacterial cell pellet.

• Mammalian Cell Samples

We recommend starting with a fresh or frozen cell pellet from 2 ml of mammalian cell culture, (e.g., from a single well of a 6-well culture plate), which should be resuspended in 100–500 µl xTractor Buffer, yielding up to 600 µl of cleared lysate). Adherent cells may be harvested by treating them with trypsin and spinning them down, or scraping them directly from the well in the presence of xTractor Buffer. Non-adherent cells may be harvested by spinning down the liquid culture.

NOTE: When lysing mammalian cells, you may substitute your standard lysis buffer for xTractor Buffer.

Clontech Laboratories, Inc.

A Takara Bio Company

1290 Terra Bella Avenue, Mountain View, CA 94043, USA

U.S. Technical Support: tech@clontech.com

United States/Canada
800.662.2566
(061515)

Asia Pacific
+1.650.919.7300

Europe
+33.(0)1.3904.6880

Japan
+81.(0)77.543.6116

IV. His-Tagged Protein Purification

1. Add 400 μ l xTractor Buffer to a spin column which has been placed in the provided collection tube, in order to equilibrate the column. Centrifuge at 11,000 x g for 1 minute at room temperature. Remove the flowthrough and discard it along with the collection tube, then place the column in a new collection tube (supplied by the user—see Section II.B).
2. Load 200–800 μ l cleared lysate (see Section III) onto the equilibrated spin column. Centrifuge at 11,000 x g for 1 minute at room temperature. Save the collection tube containing the lysate flowthrough for protein analysis and transfer the spin column to a new collection tube (supplied by the user).
3. Add 300 μ l Wash Buffer to the spin column. Centrifuge at 11,000 x g for 1 min at room temperature. Save the collection tube containing the wash flowthrough for protein analysis and transfer the spin column to a new collection tube (supplied by the user).

NOTE: Some purifications require optimization, and may benefit from addition of imidazole to the Wash Buffer. See Table 1, below, for instructions on how to prepare 1 ml of Wash Buffer containing different concentrations of imidazole (by combining different volumes of Wash Buffer and Elution Buffer).

Table 1. Adding Imidazole to Wash Buffer

Desired Imidazole Concentration	Wash Buffer Volume	Elution Buffer Volume
10 mM	980 μ l	20 μ l
20 mM	960 μ l	40 μ l
40 mM	920 μ l	80 μ l

4. Add 300 μ l Elution Buffer to the spin column. Centrifuge at 11,000 x g for 1 minute at room temperature. The collection tube should contain your eluted tagged protein, which is now ready for analysis.
- NOTE:** \geq 90% of your tagged protein can be eluted with 100 μ l of Elution Buffer.
5. Measure the amount of protein in your flowthrough samples from Steps 2 and 3, and your eluate from Step 4, using a Bradford assay or other colorimetric protein analysis method.
 6. Analyze the samples that were quantified in Step 5 using SDS-PAGE.

Contact Us For Assistance	
Customer Service/Ordering	Technical Support
Telephone: 800.662.2566 (toll-free)	Telephone: 800.662.2566 (toll-free)
Fax: 800.424.1350 (toll-free)	Fax: 650.424.1064
Web: www.clontech.com	Web: www.clontech.com
E-mail: orders@clontech.com	E-mail: tech@clontech.com

Notice to Purchaser

Our products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, *in vitro* diagnostic purposes, therapeutics, or in humans. Our products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without prior written approval of Clontech Laboratories, Inc.

Your use of this product is also subject to compliance with any applicable licensing requirements described on the product's web page at <http://www.clontech.com>. It is your responsibility to review, understand and adhere to any restrictions imposed by such statements.

Clontech® and the Clontech logo are trademarks of Clontech Laboratories, Inc. All other marks are the property of their respective owners. Certain trademarks may not be registered in all jurisdictions. Clontech is a Takara Bio Company. ©2015 Clontech Laboratories, Inc.

This document has been reviewed and approved by the Clontech Quality Assurance Department.