

SDR HyperD[®]

Solvent-detergent removal chromatography sorbent.

- Binds solvent and detergent molecules used in viral inactivation processes (TnBP and Triton* X-100).
- High recovery for proteins (exclusion limit 10 kDa).
- High sorption capacity for small hydrophobic molecules.
- Stable in acid, polar organic and oxidizing solutions.

Plasma preparations may contain viruses that are effectively removed and inactivated by combining nanofiltration (for size exclusion removal of non-lipid enveloped viruses) and treatment with non-ionic solvents and detergents (effective for lipid-coated viruses).

The elimination of solvent and detergent from biological fractions is necessary, and can be achieved by various methods including sorbent partitioning, size exclusion, affinity or batch extraction with vegetable oils combined to reverse phase on C18.


SDR HyperD is a unique sorbent designed to eliminate solvent and detergent from biological fluids.

Product description and mechanism of solvent detergent adsorption.

SDR HyperD is a composite sorbent that combines a silica-bead moiety filled with a three-dimensional cross-linked hydrophobic polymer.

The particle size distribution (40-100 µm), the small pore size of the silica beads and the nature of the polymer have been optimized for a retention of solvents and detergents typically used in viral inactivation processes (i.e. Tri-n-Butyl Phosphate – TnBP – and Triton* X-100).

Figure 1. Schematic interaction mechanism of Triton* X-100 and TnBP on SDR HyperD[®] sorbent.



Triton* X-100 interacts both with the silica surface (formation of hydrogen bonds between the silanols and the polyoxyethylene chain) and with the hydrophobic polymer moiety. TnBP interacts only with the hydrophobic polymer of the sorbent.

The adsorption mechanism involves both the silica moiety and the hydrophobic polymer. The adsorption of Triton* X-100 is proportional to the silica surface area, whereas the adsorption of TnBP is linked to binding to the organic polymer moiety. The main features of SDR HyperD sorbent are summarized in Table I and the adsorption mechanism represented in Figure 1.

Table I. Main Properties of SDR HyperD® sorbent.

Sorbent structure	Spherical silica beads filled with a three-dimensional hydrophobic polymer.
Average particle size	40-100 µm
Nature of polymer	Hydrophobic, long aliphatic chains bind solvents. 10 kDa limit prevents proteins from being retained.
Typical sample load	2-3 times the column volume with residence times of 5 min using IgG or ATIII treated solutions.
Recommended residence time	5-15 min
Binding capacity for Triton* X-100	≥90 mg/ml ⁽¹⁾
Adsorption buffer	PBS
Solvent/detergent elution buffer	PBS/Ethanol (50/50) and EtOH or/and isopropanol
Operating pH range	2-12
Sorbent pressure resistance	70 bar (1000 psi)

⁽¹⁾ Determined using 5 mg/ml Triton* X-100 in PBS, pH 7.4, 10% breakthrough, 300 cm/h.

Capacity and solvent-detergent removal efficiency.

The structure of SDR HyperD sorbent has been engineered to optimize the solvent/detergent retention. Due to the specific degree of three-dimensional polymer cross-linking, a low exclusion limit of 10 kDa allows target proteins to be "excluded" from the sorbent, and are found unrestrained in the column void volume.

On the other hand, the high specific surface area (200 m²/g) of the porous silica allows a high capacity for Triton* X-100 and TnBP. The dynamic binding capacities currently obtained are 60-80 mg/ml for Triton* X-100 and 40-50 mg/ml for TnBP at

100 cm/h (initial concentration of Triton* X-100 and TnBP in bovine plasma are respectively 10 and 5 mg/ml).

Examples of removal efficiencies from various feedstreams are summarized in Table II.

Table II. Solvent-Detergent Depletion Example.

		Before depletion	After depletion	Removal efficiency
IgG	TnBP	5,000 ppm	< 0.4 ppm	99.9%
	Triton* X-100	10,000 ppm	< 10 ppm	99.9%
ATIII	TnBP	5,000 ppm	< 0.4 ppm	99.9%
	Triton* X-100	10,000 ppm	< 10 ppm	99.9%
Bovine serum	TnBP	5,000 ppm	< 0.4 ppm	99.9%
	Triton* X-100	10,000 ppm	< 10 ppm	95.5%

Sample volume: 3.6 CV, Flow rate: 150 cm/h; Column length: 10 cm; Residence time: 4 min.

NOTE: The removal efficiency is also dependent on flow rate and column loading: i.e. when using a 10 cm column at 150 cm/h (2 CV load of bovine serum supplemented with Triton* X-100 or TnBP), a removal of 95.5% was observed for Triton* X-100; this removal efficiency was decreased to 80% when 8 CV loads were used.

Guidelines for binding capacity optimization.

As for any sorbent, the dynamic binding capacity (DBC) of SDR **HyperD** sorbent is sensitive to the linear flow rate, to the residence time on the column and to the nature of the sample. Therefore, it is recommended to start trials by loading not more than 5 CV of sample on a column with a minimum height of 15 cm, and a working flow rate not exceeding 150 cm/h. Adsorption buffer sample load and flow rate can be increased according to performance. Note that the rigid nature of SDR **HyperD** sorbent allows to use higher flow rates (i.e. >600 cm/h) for washing and cleaning, with moderate backpressure (< 2 bar).

Desorption of retained solvent/detergent.

It is typically achieved by injecting 1-10 CV of PBS/Ethanol 95° (50/50), followed by 3-10 CV of ethanol 95°. If necessary, 2-isopropanol can also be used (10 CV) in the washing sequence.

Cleaning in place and chemical stability.

SDR **HyperD** sorbent is insoluble in water and in organic solvents. It is also very stable to strong denaturing agents and chaotropic agents. The sorbent can be treated with 0.01 to 0.1 M hydrochloric acid or water-miscible organic solvent. SDR **HyperD** sorbent can also be treated with oxidizing agents such as peracetic acid (1500 ppm in sodium acetate, pH 5.0), which is a well-known bactericide and sporicide.

Recommendations for sanitization are shown in Table III.

Table III. Recommended clean-in-place method.

Alcohol/acid treatment

Wash with at least 3 CV of a solution of 20% (v/v) ethanol containing 1 M acetic acid. This solution should be injected after removal of dissolved gas at a flow rate of 10-20 cm/h (1 hour contact time). After treatment, reequilibrate with normal sterile pyrogen-free buffer.

SDR **HyperD** sorbent is supplied in a storage solution containing 20% ethanol.

For industrial applications, bulk quantities are available on request.



Ordering Information

Product	Cat. No.	Size
SDR HyperD ®	20033-065	5 mL
	20033-031	25 mL
	20033-023	100 mL
	20033-015	1 L
	20033-056	5 L
	20033-049	10 L

Reference

- Specific sorbent to remove solvent-detergent mixtures from virus-inactivated biological fluids. Guerrier, L., et al., J. Chromatogr. B, 664 (1995) 119.



Life Sciences

New York - USA

+1 516.484.5400 phone
 +1 516.801.9548 fax
 pharmafilter@pall.com

Portsmouth - Europe

+44 (0)23 9230 3303 phone
 +44 (0)23 9230 2506 fax
 BioPharmUK@europe.pall.com

Cergy - France

Pall BioSeptra
 +33 (0)1 34 20 78 00 phone
 +33 (0)1 34 20 78 78 fax
 bioseprainfo@pall.com

Visit us on the web at www.pall.com/biopharmaceutical

These products are for laboratory research use only and are not intended for human or animal diagnostic, therapeutic, or other clinical uses, unless otherwise stated. The information contained in this brochure are subject to change without notice.

Pall Corporation has offices and plants throughout the world in locations including: Argentina, Australia, Austria, Belgium, Brazil, Canada, China, France, Germany, India, Indonesia, Ireland, Italy, Japan, Korea, Malaysia, Mexico, the Netherlands, New Zealand, Norway, Poland, Puerto Rico, Russia, Singapore, South Africa, Spain, Sweden, Switzerland, Taiwan, Thailand, United Kingdom, United States and Venezuela. Distributors are located in all major industrial areas of the world.

Pall, BioSeptra, HyperD, HyperCel are trademarks of Pall Corporation.

Filtration. Separation. Solution. is a service mark of Pall Corporation. Triton is a trademark of a subsidiary of The Dow Chemical Company.

® indicates a trademark registered in the U.S.

© Pall Corporation 2005